



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

PO9031



ADULT AMBULATORY INFUSION ORDER

**Ferric Carboxymaltose  
(INJECTAFER) Infusion**

Page 1 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE**

Weight: \_\_\_\_\_ kg      Height: \_\_\_\_\_ cm

Allergies: \_\_\_\_\_

Diagnosis Code: \_\_\_\_\_

Treatment Start Date: \_\_\_\_\_ Patient to follow up with provider on date: \_\_\_\_\_

**\*\*This plan will expire after 365 days at which time a new order will need to be placed\*\***

**GUIDELINES FOR ORDERING**

1. Send **FACE SHEET and H&P or most recent chart note.**
2. **Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date: \_\_\_\_\_**

**NURSING ORDERS:**

1. TREATMENT PARAMETER – Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
2. Monitor the patient for signs and symptoms of hypersensitivity during the infusion and for at least 30 minutes after completion of the infusion. Also monitor BP following infusion.
3. Remind patient to contact provider to set up lab draw, approximately 4 weeks after treatment infusion.
4. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

**MEDICATIONS:**

**ferric carboxymaltose (INJECTAFER): (must check one)**

- Weight 50 kg or greater – 750 mg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 30 minutes
- Weight less than 50 kg – 15 mg/kg = \_\_\_\_\_ mg in sodium chloride 0.9%, intravenous, ONCE, over 30 minutes (Pharmacy to prepare in an appropriate volume)

Avoid extravasation (may cause persistent discoloration). Monitor, if extravasation occurs, discontinue administration at that site.

**Interval: (must check one)**

- 2 doses at least 7 days apart
- Other: \_\_\_\_\_

**AS NEEDED MEDICATIONS:**

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with ferric carboxymaltose



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

**Ferric Carboxymaltose  
(INJECTAFER) Infusion**

Page 2 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE**

**HYPERSENSITIVITY MEDICATIONS:**

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydramine (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

**By signing below, I represent the following:**

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in:  Oregon  \_\_\_\_\_ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

**My physician license Number is # \_\_\_\_\_ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION);** and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

**Provider signature:** \_\_\_\_\_ **Date/Time:** \_\_\_\_\_

**Printed Name:** \_\_\_\_\_ **Phone:** \_\_\_\_\_ **Fax:** \_\_\_\_\_



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

**Ferric Carboxymaltose  
(INJECTAFER) Infusion**

Page 3 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE**

***Please check the appropriate box for the patient's preferred clinic location:***

**Hillsboro Medical Center**

Infusion Services  
364 SE 8th Ave, Medical Plaza Suite 108B  
Hillsboro, OR 97123  
Phone number: (503) 681-4124  
Fax number: (503) 681-4120

**Adventist Health Portland**

Infusion Services  
10123 SE Market St  
Portland, OR 97216  
Phone number: (503) 261-6631  
Fax number: (503) 261-6756

**Mid-Columbia Medical Center**

Celilo Cancer Center  
1800 E 19th St  
The Dalles, OR 97058  
Phone number: (541) 296-7585  
Fax number: (541) 296-7610