

Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER Iron Sucrose (VENOFER) Infusion

Page 1 of 3

ACCOUNT NO. MED. REC. NO. NAME **BIRTHDATE**

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight:kg Height:cm			
Allergies:			
Diagnosis Code:			
Treatment Start Date: Patient to follow up with provider on date:			
This plan will expire after 365 days at which time a new order will need to be placed			
 GUIDELINES FOR ORDERING Send FACE SHEET and H&P or most recent chart note. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date: 			
 NURSING ORDERS: TREATMENT PARAMETER – Hold treatment and notify provider if Ferritin greater than 300 ng/mL. Instruct patient to obtain ferritin lab 30 days after infusion treatment and set up follow up appointment with provider. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes. 			
MEDICATIONS:			
iron sucrose (VENOFER): (must check one)			
 200 mg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 30 minutes, x 5 doses over 14 days 300 mg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 1.5 hours, x 3 doses (administered every 2 to 3 days) 			
AS NEEDED MEDICATIONS:			

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with iron sucrose

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health Iron Sucrose (VENOFER) Infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following: I am responsible for the care of the patient (who is I hold an active, unrestricted license to practice methat corresponds with state where you provide can	edicine in: 🛭 Oreg	gon □ (check box
state if not Oregon);	re to patient and wir	ere you are currently licensed. Specify
My physician license Number is #	be of practice and au	
Provider signature:	Dat	e/Time:
Printed Name:	Phone:	Fax:



Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU Health ADULT AMBULATORY INFUSION ORDER Iron Sucrose (VENOFER) Infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Please check the appropriate box for the patient's preferred clinic location:

☐ Hillsboro Medical Center

Infusion Services 364 SE 8th Ave, Medical Plaza Suite 108B Hillsboro, OR 97123

Phone number: (503) 681-4124 Fax number: (503) 681-4120

☐ Mid-Columbia Medical Center

Celilo Cancer Center 1800 E 19th St The Dalles, OR 97058

Phone number: (541) 296-7585 Fax number: (541) 296-7610 ☐ Adventist Health Portland

Infusion Services 10123 SE Market St Portland, OR 97216

Phone number: (503) 261-6631 Fax number: (503) 261-6756