

ADULT AMBULATORY INFUSION ORDER

Sodium Ferric Gluconate Complex
(FERRLECIT) Infusion

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

weign	t:Kg Height:cm
Allerg	es:
Diagn	osis Code:
Treatn	nent Start Date: Patient to follow up with provider on date:
This	plan will expire after 365 days at which time a new order will need to be placed
1.	Send FACE SHEET and H&P or most recent chart note. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date:
1. 2. 3. 4.	TREATMENT PARAMETERS – Hold treatment and notify provider if Ferritin greater than 300 ng/mL. Ensure patient has been scheduled for follow-up labs and visit with the provider. Monitor patient for signs and symptoms of hypotension during and following administration. Monitor patient for signs and symptoms of hypersensitivity during the infusion and for at least 30 minutes after completion of the infusion. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.
MEDI	CATIONS:
	dium ferric gluconate complex (FERRLECIT) 125 mg in sodium chloride 0.9% 100 mL, intravenous, er 1 hour
In	terval: □ Once □ Other:

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9% bolus, intravenous, 500 mL, AS NEEDED x1 dose, for vein discomfort tolerability. Give concurrently with ferric gluconate



Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Health Sodium Ferric Gluconate Complex (FERRLECIT) Infusion

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
 infusion and notify provider immediately. Administer emergency medications per the Treatment
 Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
 symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following: am responsible for the care of the patient (who is identified at the top of this form); hold an active, unrestricted license to practice medicine in: Oregon (check box					
that corresponds with state where you provide care state if not Oregon);					
My physician license Number is #(MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.					
Provider signature:	Date/Time:				
Printed Name:	_ Phone:	Fax:			
Please check the appropriate box for the patient's preferred clinic location:					
☐ Hillsboro Medical Center	☐ Adventist He				
Infusion Services	Infusion Servi				
364 SE 8th Ave, Medical Plaza Suite 108B Hillsboro, OR 97123	10123 SE Ma Portland, OR				
Phone number: (503) 681-4124	-	er: (503) 261-6631			
Fax number: (503) 681-4120	Fax number:	(503) 261-6756			
☐ Mid-Columbia Medical Center					

Phone number: (541) 296-7585 Fax number: (541) 296-7610

Celilo Cancer Center 1800 E 19th St

The Dalles, OR 97058