



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO8031



ADULT AMBULATORY INFUSION ORDER
Vedolizumab (ENTYVIO) Infusion

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
3. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.
4. Patients should not have an active ongoing infection, signs or symptoms of malignancy, or moderate to severe heart failure at the onset of therapy. Baseline liver function tests should be normal. 2. Patient should have regular monitoring for infection, malignancy, and liver abnormalities throughout therapy.

PRE-SCREENING: (Results must be available prior to initiation of therapy):

- Hepatitis B surface antigen and core antibody total test results scanned with orders.
- Tuberculin skin test or QuantiFERON Gold blood test results scanned with orders.
- Chest X-Ray result scanned with orders if TB test result is indeterminate.

LABS:

- Complete Metabolic Panel, Routine, ONCE, every visit
- CBC with differential, Routine, ONCE, every visit

NURSING ORDERS:

1. TREATMENT PARAMETER – Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
2. VITAL SIGNS – Monitor patient for signs and symptoms of hypersensitivity during the infusion.
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

MEDICATIONS:

vedolizumab (ENTYVIO) 300 mg in sodium chloride 0.9%, intravenous, ONCE over 30 minutes

Interval (must check at least one)

- Initial dosing:** on week 0, 2 and 6
- Maintenance dosing:** every 8 weeks thereafter
- Other:** _____



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Vedolizumab (ENTYVIO) Infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

AS NEEDED MEDICATIONS:

1. acetaminophen (TYLENOL) tablet, 650 mg, oral, EVERY 4 HOURS AS NEEDED for fever, headache, chills, or malaise
2. diphenhydramine (BENADRYL) capsule, 50 mg, oral, EVERY 4 HOURS AS NEEDED for itching

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydramine (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHRINE HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____



**Oregon Health & Science University
Hospital and Clinics Provider's Orders**

ADULT AMBULATORY INFUSION ORDER
Vedolizumab (ENTYVIO) Infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Please check the appropriate box for the patient's preferred clinic location:

Hillsboro Medical Center

Infusion Services
364 SE 8th Ave, Medical Plaza Suite 108B
Hillsboro, OR 97123

Phone number: (503) 681-4124

Fax number: (503) 681-4120

Adventist Health Portland

Infusion Services
10123 SE Market St
Portland, OR 97216

Phone number: (503) 261-6631

Fax number: (503) 261-6756

Mid-Columbia Medical Center

Celilo Cancer Center
1800 E 19th St
The Dalles, OR 97058

Phone number: (541) 296-7585

Fax number: (541) 296-7610